

# Exercices

## MPRI 2-6, year 2015–2016

Antoine Miné

23 September 2015

Let  $\Sigma$  be a set of states. Given a transition relation  $\tau \subseteq \Sigma \times \Sigma$ , we denote by  $\mathcal{T}[\tau]$  the set of partial finite traces obeying  $\tau$ :

$$\mathcal{T}[\tau] \stackrel{\text{def}}{=} \{(\sigma_0, \dots, \sigma_n) \in \Sigma^+ \mid \forall i < n : (\sigma_i, \sigma_{i+1}) \in \tau\} .$$

We say that the transition relation  $\tau$  generates the trace set  $\mathcal{T}[\tau]$ .

1. Give a definition for  $\mathcal{S}[T]$ , the function that, given a set of traces  $T \subseteq \Sigma^+$ , returns the smallest transition relation (for  $\subseteq$ ) that generates a set of traces containing  $T$ .
2. Prove that the pair  $\mathcal{S}$  and  $\mathcal{T}$  forms a Galois connection between trace sets in  $\mathcal{P}(\Sigma^+)$  and transition relations in  $\mathcal{P}(\Sigma \times \Sigma)$ :

$$(\mathcal{P}(\Sigma^+), \subseteq) \xleftrightarrow[\mathcal{S}]{\mathcal{T}} (\mathcal{P}(\Sigma \times \Sigma), \subseteq)$$

3. Prove that not all trace sets are generated by a transition relation. Give an example where  $\mathcal{S}$  results in an approximation.
4. Prove that the abstraction  $\mathcal{S}[T]$  does not lose any information on  $T$  if and only if  $T$  is closed at the same time by junction ( $T \hat{\ } T = T$ ), by prefix, and by suffix, and if  $\Sigma \subseteq T$ .

Consider now the set  $\mathcal{T}_\infty[\tau]$  of partial finite and infinite traces obeying  $\tau$ :

$$\mathcal{T}_\infty[\tau] \stackrel{\text{def}}{=} \{(\sigma_0, \dots, \sigma_n) \in \Sigma^+ \mid \forall i < n : (\sigma_i, \sigma_{i+1}) \in \tau\} \cup \{(\sigma_0, \dots) \in \Sigma^\omega \mid \forall i : (\sigma_i, \sigma_{i+1}) \in \tau\}$$

5. Prove that it is possible to define a new Galois connection:

$$(\mathcal{P}(\Sigma^\infty), \subseteq) \xleftrightarrow[\mathcal{S}_\infty]{\mathcal{T}_\infty} (\mathcal{P}(\Sigma \times \Sigma), \subseteq)$$

by extending the function  $\mathcal{S}$  from Question 1 to a function  $\mathcal{S}_\infty : \mathcal{P}(\Sigma^\infty) \rightarrow \mathcal{P}(\Sigma \times \Sigma)$ .

6. Prove that, for this new Galois connection,  $\mathcal{S}_\infty[T]$  can lose some information on trace sets that are closed at the same time by prefix, by suffix, and by junction and contain  $\Sigma$  (give an example).
7. Provide a necessary and sufficient condition on  $T$  such that  $\mathcal{S}_\infty[T]$  does not lose any precision in this new Galois connection.